

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number  
**WO 01/00232 A2**

- (51) International Patent Classification<sup>7</sup>: **A61K 39/00**
- (21) International Application Number: **PCT/EP00/05998**
- (22) International Filing Date: **28 June 2000 (28.06.2000)**
- (25) Filing Language: **English**
- (26) Publication Language: **English**
- (30) Priority Data:  
9915205.0 29 June 1999 (29.06.1999) GB  
0002200.4 31 January 2000 (31.01.2000) GB
- (71) Applicant (for all designated States except US):  
**SMITHKLINE BEECHAM BIOLOGICALS S.A.**  
[BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): **GARCON, Nathalie** [FR/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE). **VOSS, Gerald** [DE/BE]; SmithKline Beecham Biologicals S.A., Rue de l'Institut 89, B-1330 Rixensart (BE).
- (84) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— Without international search report and to be republished upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/00232 A2

(54) Title: **VACCINE**

(57) Abstract: A vaccine formulation for the prevention or amelioration of HIV infection in humans is provided. The vaccine comprises an HIV antigen, especially a protein which comprises Nef and/or Tat of HIV, and an immunostimulatory CpG oligonucleotide. Methods for making the vaccine formulation of the invention are described. Patients may also be treated by pre-administration of the CpG oligonucleotide prior to administration of the HIV antigen.

## VACCINE

The present invention relates to new vaccine formulations and their use in the treatment or prophylaxis of HIV infections. In particular the invention relates to the use of HIV antigen in conjunction with a CpG oligonucleotide.

HIV is the primary cause of AIDS (acquired immune deficiency syndrome) and is regarded, as one of the worlds most pressing health issues. Although there has been extensive efforts to produce a successful vaccine, efforts to-date have failed to produce one. Accordingly, there remains a need for an improved HIV immunogenic composition.

International Patent Application No. 92/06113 relates to vaccine formulation comprising the envelope protein gp160 and its naturally occurring derivative gp120 adjuvanted 3 deacylated monophosphoryl lipid A and a suitable carrier such as aluminium hydroxide.

International Patent Application No. 99/16884 describes vaccine formulations based on non-envelope HIV proteins, Nef and Tat, particularly fusions of the Nef protein with Tat, and fusions of the Nef protein with Tat or an immunological fusion partner.

Immunomodulatory oligonucleotides contain unmethylated CpG dinucleotides ("CpG") and are known (WO 96/02555, EP 468520). CpG is an abbreviation for cytosine-guanosine dinucleotide motifs present in DNA. Historically, it was observed that the DNA fraction of BCG could exert an anti-tumour effect. In further studies, synthetic oligonucleotides derived from BCG gene sequences were shown to be capable of inducing immunostimulatory effects (both in vitro and in vivo). The authors of these studies concluded that certain palindromic sequences, including a central CG motif, carried this activity. The central role of the CG motif in immunostimulation was later elucidated in a publication by Krieg, Nature 374, p546 1995. Detailed analysis has shown that the CG motif has to be in a certain

sequence context, and that such sequences are common in bacterial DNA but are rare in vertebrate DNA.

It is currently believed that this evolutionary difference allows the vertebrate  
5 immune system to detect the presence of bacterial DNA (as occurring during an infection) leading consequently to the stimulation of the immune system. The immunostimulatory sequence as defined by Krieg is:

Purine Purine CG pyrimidine pyrimidine and where the CG motif is not methylated.  
10 In certain combinations of the six nucleotides a palindromic sequence is present. Several of these motifs, either as repeats of one motif or a combination of different motifs, can be present in the same oligonucleotide. The presence of one or more of these immunostimulatory sequence containing oligonucleotides can activate various immune subsets, including natural killer cells (which produce interferon  $\gamma$  and have  
15 cytolytic activity) and macrophages (Wooldrige et al Vol 89 (no. 8), 1977). Although other unmethylated CpG containing sequences not having this consensus sequence have now been shown to be immunomodulatory.

The present invention provides an improved vaccine formulation comprising a CpG  
20 oligonucleotide and an HIV antigen selected from the group HIV gp160, HIV gp120; an HIV Nef protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Tat protein or derivative thereof; or an HIV Tat protein or derivative thereof linked to either (i) a fusion protein or (ii) an HIV Nef protein or derivative thereof; or an HIV Nef protein or derivative thereof linked to an HIV Tat protein or  
25 derivative thereof and a fusion partner.

By 'fusion partner' is meant any protein sequence that is not Tat or Nef. Preferably the fusion partner is protein D or its lipidated derivative Lipoprotein D, from *Haemophilus influenzae* B. In particular, it is preferred that the N-terminal third,  
30 i.e. approximately the first 100-130 amino acids are utilised. This is represented herein as Lipo D 1/3. In a preferred embodiment of the invention the Nef protein

or derivative thereof may be linked to the Tat protein or derivative thereof. Such Nef-Tat fusions may optionally also be linked to a fusion partner, such as protein D.

If present, the fusion partner is normally linked to the N-terminus of the Nef or Tat  
5 protein.

Derivatives encompassed within the present invention include molecules with a C terminal Histidine tail, which preferably comprises between 5-10 Histidine residues.

Generally, a histidine tail containing n residues is represented herein as His (n).

10 The presence of a histidine (or 'His') tail aids purification. More specifically, the invention provides vaccines comprising proteins with the following structure:

	Lipo D 1/3	-	Nef	-	His (6)
15	Lipo D 1/3	-	Nef-Tat	-	His (6)
	Prot D 1/3	-	Nef	-	His (6)
	Prot D 1/3	-	Nef-Tat	-	His (6)
20			Nef-Tat		His (6)

In a preferred embodiment the proteins are expressed with a Histidine tail comprising between 5 to 10 and preferably six Histidine residues. These are  
25 advantageous in aiding purification. Separate expression, in yeast (*Saccharomyces cerevisiae*), of Nef (Macreadie IG et al 1993, *Yeast* 9 (6) 565-573) and Tat (Braddock M et al, 1989, *Cell* 58 (2) 269-79) has already been reported. Nef protein only is myristilated. The present invention also provides vaccines comprising Nef and/or Tat separately. The DNA and amino acid sequences of  
30 representative Nef-His, Tat-His and of Nef-Tat-His fusion proteins as set forth in WO99/16884.

Particularly preferred vaccines according to the invention comprise an HIV fusion protein such as Nef-Tat and optionally an additional HIV protein. Particularly preferred in a vaccine formulation according to the invention is a combination of an HIV fusion protein plus gp120, more particularly Nef-Tat with gp120.

5

Derivatives encompassed within the present invention also include mutated proteins. The term 'mutated' is used herein to mean a molecule which has undergone deletion, addition or substitution of one or more amino acids using well known techniques for site directed mutagenesis or any other conventional method.

10

Vaccine preparation is generally described in Vaccine Design - The subunit and adjuvant approach (Ed. Powell and Newman) Pharmaceutical Biotechnology Vol. 6 Plenum Press 1995. Encapsulation within liposomes is described by Fullerton, US Patent 4,235,877.

15

The preferred oligonucleotides preferably contain two or more CpG motifs separated by six or more nucleotides. The oligonucleotides of the present invention are typically deoxynucleotides. In a preferred embodiment the internucleotide in the oligonucleotide is phosphorodithioate, or more preferably a phosphorodithioate bond, although phosphodiester and other internucleotide bonds are within the scope of the invention including oligonucleotides with mixed internucleotide linkages.

20

Preferred oligonucleotides have the following sequences:

Oligo (internal designation*)	5'-SEQUENCE-3'	CpG	Thio
WD1001	TCC ATG ACG TTC CTG ACG TT	+	+
WD1002	TCT CCC AGC GTG CGC CAT	+	+
WD1003	ACC GAT AAC GTT GCC GGT GAC G	+	-
WD1004	G*G*G GTC AAC GTT GAG* G*G*G* G*G	+	Mix
WD1005	TCC ATG AGC TTC CTG AGC TT	-	+

WD1006	TCC ATG ACG TTC CTG ACG TT	+	-
WD1007	ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG	+	+
	TCG TCG TTT TGT CGT TTT GTC GTT	+	+

\* alternatively referred to as WD001-WD007

In the above table a + in the Thio column indicates the presence of a thioate modification. 'Mix' indicates a mixture of thioate modification and sequence without thioate modification (the asterisks indicate the linkages with a thioate modification). A - in the Thio column indicates absence of a thioate modification. A + in the CpG column indicates the presence of a CpG motif and a - in the CpG column indicates absence of a CpG motif. For example WD1005 contains a GpC rather than a CpG motif, thus it is marked with a - in the CpG column of the table. WD1007 contains a palindromic motif (GACGTC) as well as other non-palindromic CpG sequences. This is also within the scope of a CpG oligonucleotide as the term is used in the present application.

The CpG oligonucleotides utilised in the present invention may be synthesized by any method known in the art (eg EP 0 468 520). Conveniently, such oligonucleotides may be synthesized utilising an automated synthesizer. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are described in US 5,666,153, US 5,278,302 and WO 95/26204.

The amount of protein in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccinees. Such amount will vary depending upon which specific immunogen is employed and how it is presented. Generally, it is expected that each dose will comprise 1-1000 µg of protein, preferably 2-500 µg, most preferably 5-250 µg. An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of appropriate immune responses in subjects. Following an initial vaccination, subjects may receive one or several booster immunisations adequately spaced.

It is also possible to pre-administer the CpG oligonucleotide shortly prior to vaccination with the HIV antigen, for example 1 day before.

Accordingly, according to another aspect of the invention there is provided a  
5 method of treating a human in need thereof either with a vaccine composition comprising an HIV antigen and a CpG oligonucleotide (as hereinabove defined) or with the CpG oligonucleotide followed after a suitable time by the HIV antigen.

There is also provided a kit comprising effective amounts of a CpG oligonucleotide-  
10 containing formulation for use as a priming formulation for pre-administration to human patients and an HIV antigen for injection at some suitable time later, as described hereinabove.

Preferred CpG oligonucleotides are those indicated in the table hereinabove.  
15

Suitably the CpG will be present in the range 10 µg per dose to 2000 µg, preferably 50-1000 µg, such as about 50 or about 500 or about 1000 µg per dose.

Suitably the vaccine used in the present invention may comprise a carrier such as an  
20 aluminium salt, eg aluminium hydroxide (Al(OH)<sub>3</sub>), aluminium phosphate or aluminium phosphate sulfate (alum), or a non-toxic oil in water emulsion or a mixture thereof.

If an aluminium salt (preferably aluminium hydroxide) is used as a carrier it is  
25 generally present in the range of 50 to 100 µg, preferably 100 to 500 µg per dose.

Non-toxic oil in water emulsions preferably contain a non-toxic oil, eg squalene and an emulsifier such as (polysorbitan monoleate) Tween 80, in an aqueous carrier such as phosphate buffered saline.  
30

If desired the vaccine used in the present invention may comprise an additional adjuvant, preferably a saponin adjuvant such as QS21 as described for example in

WO 9517210, optionally in the presence of a sterol, such as cholesterol as described for example in PCT/EP96/01464. The vaccine of the invention may also comprise monophosphoryl lipid A and derivatives thereof known in the art. A preferred derivative is 3 de-O-acylated monophosphoryl lipid A, described in British Patent

5 No. GB 2 220 211.

Accordingly vaccine formulations of the present invention may additionally comprise other pharmaceutical excipients or immunostimulants. In a preferred embodiment the vaccine formulation additionally comprises an aluminium salt,

10 preferably aluminium hydroxide. In a further embodiment a saponin adjuvant may also be included such as QS21 (Aquila).

The present invention will now be described with reference to the following examples:



## EXAMPLES

### Example 1 - IMMUNOGENICITY STUDIES USING HIV-1 GP120

#### 5 FORMULATED WITH CPG OR CPG/ALUM

#### Evaluation of CpG and CpG/alum in rhesus monkeys

##### *Experiment outline*

An immunogenicity study was conducted to evaluate the adjuvant effect of CpG in non-human primates. Groups of five monkeys were immunized twice with HIV-1<sub>W6.1D</sub> gp120 in combination with CpG or CpG/alum. After the second immunization the immune response of the animals was assessed. Antibodies to gp120 and lymphoproliferative as well as cytokine responses were evaluated.

#### Groups of monkeys

15

Group	antigen	Adjuvant
1	HIV-1 <sub>W6.1</sub> gp120	CpG/alum
2	HIV-1 <sub>W6.1</sub> gp120	CpG

#### Formulation

##### *Component batches used:*

20

COMPONENT	BRAND	BATCH NUMBER	CONCENTRATION (MG/ML)	BUFFER
Gp120*		47/025	0.456	PBS
Al(OH) <sub>3</sub>	Superfos	96A0089	10.380	H <sub>2</sub> O
CpG		WD001	5	H <sub>2</sub> O

\* gp120 of HIV-1 clone W6.1D (Groenink et al., 1991)

**Formulation process:**

Formulations were prepared one day before each injection. All incubations were carried out at room temperature with agitation.

5

**CpG/alum group 1 (500µl/dose)**

Gp120 (50µg) was adsorbed on 500µg of Al(OH)<sub>3</sub> for 1 hour. The formulation was buffered with a 10-fold concentrated PO<sub>4</sub>/NaCl pH 6.8 solution before addition of  
10 500µg of WD001. After 15 min, 50 µg/ml of thiomersal was added as preservative.

**CpG group 2 (500µl/dose)**

Gp120 (50µg) was diluted in PO<sub>4</sub>/NaCl buffer pH 6.8 before addition of 500µg of  
15 CpG WD001. After 15 min, 50 µg/ml of thiomersal was added as preservative.

**Immunological methods**

Five rhesus monkeys (*Macaca mulatta*). per group were immunized twice  
20 intramuscularly with 500 µl of vaccine at a four-week-interval. Sera and peripheral blood mononuclear cells (PBMC) were taken at several occasions.

Gp120-specific antibodies in sera of immunized monkeys were determined in an ELISA as described in Mooij et al., 1998. Briefly, ELISA plates were coated with 1  
25 µg/ml gp120, saturated with BSA and subjected to incubation with rhesus monkey sera followed by α-human biotinylated antibody. Finally, HRP-streptavidin was added and quantified in a color reaction using OPD as substrate.

Lymphoproliferation was assessed by using density gradient-purified PBMC from  
30 immunized rhesus monkeys. Cells were seeded in quadruplicates at 1x10<sup>5</sup> in 100 µl RPMI/5 % FCS per well in round bottom 96 well plates. Then another 100 µl of

medium alone or containing soluble gp120 (10  $\mu$ g/ml) were added and parallel cultures were incubated for 48 hrs. Thereafter, 100  $\mu$ l culture supernatant were replaced by fresh medium containing 1  $\mu$ Ci [ $^3$ H]-thymidine. After 16 hrs cells were harvested onto filter plates and incorporated radioactivity was determined in a  $\beta$ -counter. Results are expressed in cpm and in stimulation indices (SI, = cpm antigen-containing cultures/cpm medium alone cultures), SI greater than 3 are considered as a positive response.

Flat bottom 96 well plates were prepared by coating an IFN- $\gamma$ -specific capture antibody in 50  $\mu$ l PBS for 4 hrs at 37 °C. The plates were washed three times and PBMC were seeded similar to lymphoproliferation assays. After 48 hrs of culture the plates were washed thrice with PBS/0.05 % Tween 20 and 50  $\mu$ l of biotinylated secondary IFN- $\gamma$ -specific antibody diluted in PBS/Tween/1 % FCS were added for 2 hrs. The plates were washed again and a gold-conjugated  $\alpha$ -biotin antibody was incubated for 1 hr. After additional washings the ELIspots were visualized by using a silver enhancing kit (50  $\mu$ l per well). The reaction was stopped after approx. 30 min by adding deionized water. Cytokine-secreting cells were enumerated by microscopic examination.

## 20 Results

Analysis of gp120-specific antibodies in sera of the monkeys revealed that all animals in the two groups had acquired specific immune responses (Table 1 and Figure 1). Some low responses were detectable in two animals already after one immunization. After the second immunization all animals exhibited high titers of gp120-specific antibodies.

Table 1

		d - 12	average	14 d post I	average	7 d post II	average	28 d post II	average
Group 1	N5V	<100		188		6445		2014	
	1055	<100		50		4699		1484	
	R441	<100	<100	50	<50	2942	5530	827	2504
	BXJ	<100		50		5808		4013	
	CPJ	<100		50		7758		4184	
Group 2	NJ5	<100		50		734		729	
	X006	<100		50		4182		1171	
	1041	<100	<100	50	<50	2834	3358	400	963
	R361	<100		258		5227		1821	
	BBE	<100		50		3811		693	

- 5 Induction of specific lymphoproliferation by immunization with gp120 in combination with CpG or CpG/alum was evaluated before immunization and 6 days post secondary immunization. All 10 animals did not exhibit any specific lymphoproliferation ( $SI > 3$ ) at the study start (data not shown). In contrast, all animals in group 1 possessed strong lymphoproliferative responses 6 days post boost
- 10 immunization (Figure 2). Similarly, all except one animal in group 2 showed specific lymphoproliferation.

- The presence of gp120-specific IFN- $\gamma$ -secreting cells was investigated in all monkeys before immunization and 6 days after the second dose. IFN- $\gamma$ -secreting
- 15 cells could not be evaluated from pre-immunization samples due to technical difficulties. However, such cells were detectable after secondary immunization (Figure 3). All animals in both groups exhibited an antigen-specific response.

### Conclusions


- 20 Immunization with HIV-1 gp120 in combination with CpG or CpG/alum induces immune responses in non-human primates. These responses include gp120-specific antibodies, lymphoproliferation and IFN- $\gamma$ -secreting cells.

**Example 2 - NEF-TAT AND MUTATED NEF-TAT FORMULATION**

The antigens prepared according to WO99/16884 were dialysed against NaCl 150mM to eliminate the  $\text{PO}_4$  ions that inhibit the adsorption of gp120 on  $\text{Al}(\text{OH})_3$ .

5

The formulation sequence is as described below:



- 20 The antigens were incubated with the CpG before adsorption on  $\text{Al}(\text{OH})_3$  to favour a potential interaction between the His tail of the Nef-Tat and Nef antigens and the oligonucleotide.

***Formulation Characterisation***

- 25 The formulation was centrifuged and the supernatant was analysed by SDS-PAGE (Novex 4-20%, reducing conditions) to check the adsorption of HIV Nef-Tat. The gel was stained with silver. Gel analysis of the formulation revealed that most of the Nef or Nef-Tat protein was adsorbed onto alum.

30

**Example 3 - IMMUNIZATION AND SHIV CHALLENGE EXPERIMENT IN RHESUS MONKEYS.*****Vaccine preparation***

- 5 A vaccine formulation was prepared in accordance with the invention comprising a Nef-Tat fusion protein and gp120 together with an oligonucleotide containing unmethylated CpG dinucleotide motifs and aluminium hydroxide as carrier.

Preparation of CpG oligonucleotide solution: CpG dry powder is dissolved in H<sub>2</sub>O to  
10 give a solution of 5 mg/ml CpG.

***Preparation of CpG formulation.***

The 3 antigens were dialyzed against NaCl 150 mM to eliminate the phosphate ions that inhibit the adsorption of gp120 on aluminium hydroxide.

- 15 The antigens diluted in H<sub>2</sub>O (100 µg gp120, 20 µg NefTat and 20 µg SIV Nef) were incubated with the CpG solution (500 µg CpG) for 30 min before adsorption on Al(OH)<sub>3</sub> to favor a potential interaction between the His tail of NefTat and Nef antigens and the oligonucleotide (stronger immunostimulatory effect of CpG described when bound to the antigen compared to free CpG). Then Al(OH)<sub>3</sub> (500  
20 µg), 10 fold concentrated NaCl and 1 µg/ml thiomersal as preservative were consecutively added at 5 min intervals.

All incubations were carried out at room temperature with agitation.

25 ***Study***

Four rhesus monkeys were immunized intramuscularly at 0, 1 and 3 months with the vaccine composition.

One month after the last immunization all animals were challenged with a pathogenic SHIV (strain 89.6p). From the week of challenge (wk16) blood samples were taken periodically at the indicated time points to determine the % of CD4-positive cells among peripheral blood mononuclear cells by FACS analysis (Figure 4) and the  
5 concentration of RNA viral genomes in the plasma by bDNA assay (Figure 5).

### ***Results***

All animals became infected after challenge with SHIV<sub>89.6p</sub>.

- 10 CD4-positive cells declined after challenge. All four animals exhibited a slight decrease in CD4-positive cells and recovered to baseline levels over time (Figure 4). In contrast, three out of four animals in an adjuvant alone control group exhibited a drop of CD4-positive cells below 5 % and never recovered.
- 15 Virus load data are almost the inverse of CD4 data (Figure 5). Virus load decreased in the animals that had received NefTat + gp120 formulated with CpG containing oligonucleotide, but remained high in three out of four control animals which received only an adjuvant.
- 20 At week 44 2/4 animals in a control group treated with only an adjuvant had to be euthanized due to AIDS-like symptoms.

### ***Conclusions***

- The combination of gp120 and NefTat (in the presence of SIV Nef) with CpG  
25 adjuvant prevents the loss of CD4-positive cells and reduces the virus load in animals infected with pathogenic SHIV<sub>89.6p</sub>.

**Figure legends**

Figure 1: HIV-1 gp120-specific antibody responses in immunized rhesus monkeys. Specific antibodies were evaluated in an ELISA. Individual values from multiple  
5 time points for each animal are shown in Table 1, and group averages are shown in Table 1 and as a graphic in figure 1.

Figure 2: Gp120-specific lymphoproliferation in immunized rhesus monkeys 6 days post second immunization. PBMC were stimulated with gp120 antigen and  
10 lymphoproliferative responses were measured by <sup>3</sup>H-thymidine incorporation. Results are expressed in cpm and as SI.

Figure 3: Gp120-specific IFN- $\gamma$ -secreting cells from immunized rhesus monkeys. IFN- $\gamma$ -secreting cells were visualized by the ELISpot method. Cytokine-secreting  
15 cells resulting in a colored spot were enumerated by microscopic examination and results are expressed semi-quantitatively (- = 0-5, + = 5-15, ++ = 15-35, +++ = 35-50, ++++ = >50).

Figures 4 and 5: CD4 cell count and viral load data from Example 3 in which  
20 rhesus monkeys were immunised with Nef-Tat + gp120 formulated with CpG and challenged with pathogenic SHIV.

**References**

Groenink, M., Fouchier, R.A., De Goede, R.E., et al. (1991). Phenotypic  
25 heterogeneity in a panel of infectious molecular human immunodeficiency virus type 1 clones derived from a single individual. *J. Virol.* 65:1968-1975.

Mooij, P., Van der Kolk, M., Bogers, M.J.M., et al. (1998). A clinically  
relevant HIV-1 subunit vaccine protects rhesus macaques from in vivo passaged  
30 simian-human immunodeficiency virus infection. *AIDS* 12:F15-F22.



### Claims

1. A vaccine formulation comprising a HIV antigen and an immunostimulatory CpG oligonucleotide.
- 5 2. A vaccine as claimed in claim 1 wherein the antigen is selected from the group, gp120, gp160, Nef, Tat, Nef-Tat fusions or immunologically equivalent derivatives.
3. A vaccine as claimed in claim 1 or 2 wherein the antigen is fused to Protein D,  
10 or lipoprotein D or a fragment thereof, from Haemophilus influenzae.
4. A vaccine as claimed in any one of claims 1 to 3 wherein the antigen comprises a Nef-Tat fusion protein
- 15 5. A vaccine as claimed in claim 4 further comprising HIV gp120.
6. A vaccine formulation as claimed herein additionally comprising an aluminium salt or a saponin adjuvant.
- 20 7. A vaccine as claimed herein wherein the oligonucleotide comprises two CpG dinucleotides.
8. A vaccine as claimed herein wherein the CpG oligonucleotide is between 15-45 nucleotides in length.
- 25 9. A vaccine as claimed herein wherein the CpG oligonucleotide comprises at least one phosphorothioate internucleotide bond.
10. A vaccine as claimed herein wherein the oligonucleotide is selected from the  
30 group:

Oligo (internal designation)	5'-SEQUENCE-3'	CpG	Thio
WD1001	TCC ATG ACG TTC CTG ACG TT	+	+
WD1002	TCT CCC AGC GTG CGC CAT	+	+
WD1003	ACC GAT AAC GTT GCC GGT GAC G	+	-
WD1004	G*G*G GTC AAC GTT GAG* G*G*G* G*G	+	Mix
WD1005	TCC ATG AGC TTC CTG AGC TT	-	+
WD1006	TCC ATG ACG TTC CTG ACG TT	+	-
WD1007	ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG	+	+
	TCG TCG TTT TGT CGT TTT GTC GTT	+	+

11. A method for the prevention or amelioration of HIV infection a patient, comprising administering an effective amount of a vaccine of claims 1 to 10 to patient.

5


12. A vaccine as claimed herein for use as a medicament.

13. A method of producing a vaccine as claimed in any of claim 1 to 8 comprising admixing an HIV antigen and a CpG immunostimulatory oligonucleotide.

10

14. A method for the prevention or amelioration of HIV infection in a patient, comprising administering an effective amount of a CpG oligonucleotide followed after a suitable time by an effective amount of an HIV antigen.

1/5

**Fig. 1**

2/5




Fig. 2b




Fig. 2d




Fig. 2a




Fig. 2c

Fig. 2

3/5

Group 1						
	N5V	1055	R441	BXJ	CPJ	
gp120	+	++	++	+	+++	5/5
Unstimulated cells	-	-	-	-	-	

Group 2						
	NJ5	X006	1041	R361	BBE	
gp120	+++	+++	++	+++	+	5/5
Unstimulated cells	-	+	+	+	-	

Fig. 3

4/5




Fig. 4

5/5




Fig. 5

THOMSON

DELPHION

RESEARCH

PRODUCTS

INSIDE DELPHION

My Account | Products






Search: Quick Number Boolean Advanced Derwent

dawn\_rhoads:  
KLAR000  
Unlimited Subscriber

## Order Accepted, Thank You

Your order number **8887128** has been accepted.Order Client Reference: **4239-66899**

- » Shopping Cart
- » Order Form
- » Order Status
- » Subscription Status
- » Registration Information
- » Change Password
- » Preferences

Download /Recipient	Product	Description	Count	Units	Unit Price (USD)
 Click to download	PDF document (download)	EP00092574B1: OLIGONUCLEOTIDE THERAPEUTIC AGENT AND METHODS OF MAKING SAME	14	1	\$0.00
 Click to download	PDF document (download)	W000100232A3: USE OF CPG AS AN ADJUVANT FOR HIV VACCINE	4	1	\$0.00
 Click to download	PDF document (download)	W000100232A2: VACCINE	23	1	\$0.00
 Click to download	PDF document (download)	W009526204A1: IMMUNE STIMULATION BY PHOSPHOROTHIOATE OLIGONUCLEOTIDE ANALOGS	37	1	\$0.00
 Click to download	Core collections PDF (external source)	EP011198249A2: USE OF CPG AS AN ADJUVANT FOR HIV VACCINE W000100232A3 delivered as substitute.	4	1	\$0.00
<b>Total Price:</b>					

[View Printable Receipt](#)

Order placed on 2004-01-15 at 17:03:11. Account: dawn\_rhoads



To download each PDF image, click on this icon in the "Recipient" column. If you prefer to save the file directly to disk, use your right mouse button to click the icon (this shows a pop-up menu) and select "Save Link As..." or "Save Target As...". When the download is complete, you'll need to launch Adobe Acrobat to open the file.

If you have any problems receiving your order, please [contact us](#), [email us](#).

To view all your completed orders, click the Order Status button.





## The Delphion Integrated View

Buy Now: [More choices...](#)

Tools: Add to Work File: [Create new Work File](#)

View: [Expand Details](#) | [INPADOC](#) | Jump to:  ☐ Go to: [Derwent...](#)

☐ [Email](#)

🔍 Title: **EP1198249A2: USE OF CPG AS AN ADJUVANT FOR HIV VACCINE**

🔍 Country: **EP** European Patent Office (EPO)  
 🔍 Kind: **A2** Publ. of Application without search report

🔍 Inventor: **GARCON, Nathalie;**  
**VOSS, Gerald;**

(Try)

🔍 Assignee: **SmithKline Beecham Biologics SA**  
[News, Profiles, Stocks and More about this company](#)

🔍 Published / Filed: **April 24, 2002 / June 28, 2000**

🔍 Application Number: **EP2000000943919**

🔍 IPC Code: **A61K 39/39; A61K 39/21; A61P 33/06; A61K 39/39; A61K 39/21;**

🔍 Priority Number: **June 29, 1999 GB1999000015205**  
**Jan. 31, 2000 GB2000000002200**

🔍 Attorney, Agent or Firm: **Tyrrell, Arthur William Russell, Dr. ;**

🔍 INPADOC [Show legal status actions](#)

Legal Status: **AL AT BE CH CY DE DK ES FI FR GB GR IE IT LI LT LU LV MC MK NL**  
 🔍 Designated Country: **RO SI**

🔍 Family: [Show 8 known family members](#)

🔍 Other Abstract Info: **CHEMABS 134(07)085122Z DERABS C2001-122974 DERABS C2001-476172**



[Nominate](#)

[this for the Gallery...](#)

© 1997-2002 Delphion, Inc. [Research Subscriptions](#) | [Privacy Policy](#) | [Terms & Conditions](#) | [Site Map](#) | [Contact Us](#)

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number  
**WO 01/00232 A3**

(51) International Patent Classification<sup>7</sup>: **A61K 39/39, 39/21, A61P 33/06 // (A61K 39/39, 39:21)**

(21) International Application Number: **PCT/EP00/05998**

(22) International Filing Date: **28 June 2000 (28.06.2000)**

(25) Filing Language: **English**

(26) Publication Language: **English**

(30) Priority Data:  
9915205.0 29 June 1999 (29.06.1999) GB  
0002200.4 31 January 2000 (31.01.2000) GB

(71) Applicant (for all designated States except US):  
**SMITHKLINE BEECHAM BIOLOGICALS S.A.**  
[BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **GARCON, Nathalie**  
[FR/BE]; SmithKline Beecham Biologicals S.A., Rue de  
l'Institut 89, B-1330 Rixensart (BE). **VOSS, Gerald**  
[DE/BE]; SmithKline Beecham Biologicals S.A., Rue de  
l'Institut 89, B-1330 Rixensart (BE).

(74) Agent: **PRIVETT, Kathryn**; SmithKline Beecham,  
Corporate Intellectual Property, Two New Horizons Court,  
Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SI, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

**Published:**

- With international search report.
- Before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments.

(88) Date of publication of the international search report:  
**25 May 2001**

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 01/00232 A3

(54) Title: **USE OF CPG AS AN ADJUVANT FOR HIV VACCINE**

(57) Abstract: A vaccine formulation for the prevention or amelioration of HIV infection in humans is provided. The vaccine comprises an HIV antigen, especially a protein which comprises Nef and/or Tat of HIV, and an immunostimulatory CpG oligonucleotide. Methods for making the vaccine formulation of the invention are described. Patients may also be treated by pre-administration of the CpG oligonucleotide prior to administration of the HIV antigen.

# INTERNATIONAL SEARCH REPORT

International Application No  
PC1/EP 00/05998

A. CLASSIFICATION OF SUBJECT MATTER  
IPC 7 A61K39/39 A61K39/21 A61P33/06 //(A61K39/39,39:21)

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 7 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, BIOSIS, WPI Data, MEDLINE, CHEM ABS Data, EMBASE, SCISEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	DEML LUDWIG ET AL: "Immunostimulatory CpG motifs trigger a T helper-1 immune response to human immunodeficiency virus type-1 (HIV-1) gp160 envelope proteins." CLINICAL CHEMISTRY AND LABORATORY MEDICINE, vol. 37, no. 3, March 1999 (1999-03), pages 199-204, XP000982148 ISSN: 1434-6621 the whole document	1-14
X	WO 99 27961 A (POWDERJECT VACCINES INC) 10 June 1999 (1999-06-10) abstract page 12, line 6 - line 29 -/-	1-14

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents:

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*G\* document member of the same patent family

Date of the actual completion of the international search

8 March 2001

Date of mailing of the international search report

19/03/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040. Tx. 31 651 epo nl.  
Fax: (+31-70) 340-3016

Authorized officer

Niemann, F

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/EP 00/05998

## C. (Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 99 16884 A (GODART STEPHANE ANDRE GEORGES ; SMITHKLINE BEECHAM BIOLOG (BE); BRU) 8 April 1999 (1999-04-08) cited in the application the whole document ---	
A	WO 92 06113 A (SMITHKLINE BEECHAM BIOLOG) 16 April 1992 (1992-04-16) cited in the application the whole document ---	
A	WO 96 02555 A (UNIV IOWA RES FOUND) 1 February 1996 (1996-02-01) cited in the application the whole document ---	
A	WO 98 37919 A (UNIV IOWA RES FOUND) 3 September 1998 (1998-09-03) the whole document ---	
P, X	WO 99 33488 A (DALEMANS WILFRIED L J ; PRIEELS JEAN PAUL (BE); SMITHKLINE BEECHAM) 8 July 1999 (1999-07-08) claims 1-8 ---	1,6-14
P, X	WO 00 23105 A (SMITHKLINE BEECHAM BIOLOG ; GARCON NATHALIE (BE)) 27 April 2000 (2000-04-27) claims 1-26 ---	1,6-14
E	WO 00 62800 A (SMITHKLINE BEECHAM BIOLOG ; FRIEDE MARTIN (BE); GARCON NATHALIE (BE)) 26 October 2000 (2000-10-26) claims 1-29 ---	1,6-14
E	WO 00 67787 A (MOSS RONALD B ; IMMUNE RESPONSE CORP INC (US)) 16 November 2000 (2000-11-16) claims 1-36 -----	1,6-14

# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PC1/EP 00/05998

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9927961 A	10-06-1999	AU 1619999 A EP 1035867 A	16-06-1999 20-09-2000
WO 9916884 A	08-04-1999	AU 1025599 A BR 9812547 A CN 1279718 T EP 1015596 A NO 20001508 A PL 339432 A	23-04-1999 25-07-2000 10-01-2001 05-07-2000 18-05-2000 18-12-2000
WO 9206113 A	16-04-1992	AU 654970 B AU 8510991 A CA 2092827 A EP 0550485 A EP 0644201 A IE 913385 A JP 6501151 T MX 9101275 A NZ 239948 A PT 99063 A,B ZA 9107747 A AP 368 A BR 9107294 A CN 1064891 A CZ 9301957 A WO 9216556 A FI 934133 A HU 67011 A NO 933342 A SK 82593 A	01-12-1994 28-04-1992 29-03-1992 14-07-1993 22-03-1995 08-04-1992 10-02-1994 04-05-1992 27-07-1993 31-08-1992 30-09-1992 28-10-1994 14-06-1994 30-09-1992 18-05-1994 01-10-1992 21-09-1993 30-01-1995 20-09-1993 12-01-1994
WO 9602555 A	01-02-1996	AU 713040 B AU 1912795 A CA 2194761 A EP 0772619 A JP 10506265 T US 6008200 A	18-11-1999 16-02-1996 01-02-1996 14-05-1997 23-06-1998 28-12-1999
WO 9837919 A	03-09-1998	AU 6667498 A EP 1039935 A	18-09-1998 04-10-2000
WO 9933488 A	08-07-1999	AU 2419099 A AU 729336 B AU 2419199 A BR 9814483 A BR 9814487 A WO 9933868 A EP 1039930 A EP 1040123 A NO 20003302 A NO 20003303 A	19-07-1999 01-02-2001 19-07-1999 10-10-2000 10-10-2000 08-07-1999 04-10-2000 04-10-2000 18-08-2000 04-08-2000
WO 0023105 A	27-04-2000	AU 1151800 A	08-05-2000
WO 0062800 A	26-10-2000	AU 4114900 A	02-11-2000
WO 0067787 A	16-11-2000	AU 4992900 A	21-11-2000

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
4 January 2001 (04.01.2001)

PCT

(10) International Publication Number  
**WO 01/00232 A2**

- (51) International Patent Classification<sup>7</sup>: A61K 39/00
- (21) International Application Number: PCT/EP00/05998
- (22) International Filing Date: 28 June 2000 (28.06.2000)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
9915205.0 29 June 1999 (29.06.1999) GB  
0002200.4 31 January 2000 (31.01.2000) GB
- (71) Applicant (*for all designated States except US*):  
SMITHKLINE BEECHAM BIOLOGICALS S.A.  
[BE/BE]; Rue de l'Institut 89, B-1330 Rixensart (BE).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): GARCON, Nathalie  
[FR/BE]; SmithKline Beecham Biologicals S.A., Rue de  
l'Institut 89, B-1330 Rixensart (BE). VOSS, Gerald  
[DE/BE]; SmithKline Beecham Biologicals S.A., Rue de  
l'Institut 89, B-1330 Rixensart (BE).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ,  
DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,  
HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ,  
NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM,  
TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).
- Published:**  
— Without international search report and to be republished  
upon receipt of that report.
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*



WO 01/00232 A2

(54) Title: VACCINE

(57) Abstract: A vaccine formulation for the prevention or amelioration of HIV infection in humans is provided. The vaccine comprises an HIV antigen, especially a protein which comprises Nef and/or Tat of HIV, and an immunostimulatory CpG oligonucleotide. Methods for making the vaccine formulation of the invention are described. Patients may also be treated by pre-administration of the CpG oligonucleotide prior to administration of the HIV antigen.

## VACCINE

The present invention relates to new vaccine formulations and their use in the treatment or prophylaxis of HIV infections. In particular the invention relates to the use of HIV antigen in conjunction with a CpG oligonucleotide.

HIV is the primary cause of AIDS (acquired immune deficiency syndrome) and is regarded, as one of the worlds most pressing health issues. Although there has been extensive efforts to produce a successful vaccine, efforts to-date have failed to produce one. Accordingly, there remains a need for an improved HIV immunogenic composition.

International Patent Application No. 92/06113 relates to vaccine formulation comprising the envelope protein gp160 and its naturally occurring derivative gp120 adjuvanted 3 deacylated monophosphoryl lipid A and a suitable carrier such as aluminium hydroxide.

International Patent Application No. 99/16884 describes vaccine formulations based on non-envelope HIV proteins, Nef and Tat, particularly fusions of the Nef protein with Tat, and fusions of the Nef protein with Tat or an immunological fusion partner.

Immunomodulatory oligonucleotides contain unmethylated CpG dinucleotides ("CpG") and are known (WO 96/02555, EP 468520). CpG is an abbreviation for cytosine-guanosine dinucleotide motifs present in DNA. Historically, it was observed that the DNA fraction of BCG could exert an anti-tumour effect. In further studies, synthetic oligonucleotides derived from BCG gene sequences were shown to be capable of inducing immunostimulatory effects (both in vitro and in vivo). The authors of these studies concluded that certain palindromic sequences, including a central CG motif, carried this activity. The central role of the CG motif in immunostimulation was later elucidated in a publication by Krieg, Nature 374, p546 1995. Detailed analysis has shown that the CG motif has to be in a certain

sequence context, and that such sequences are common in bacterial DNA but are rare in vertebrate DNA.

It is currently believed that this evolutionary difference allows the vertebrate  
5 immune system to detect the presence of bacterial DNA (as occurring during an infection) leading consequently to the stimulation of the immune system. The immunostimulatory sequence as defined by Krieg is:

Purine Purine CG pyrimidine pyrimidine and where the CG motif is not methylated.  
10 In certain combinations of the six nucleotides a palindromic sequence is present. Several of these motifs, either as repeats of one motif or a combination of different motifs, can be present in the same oligonucleotide. The presence of one or more of these immunostimulatory sequence containing oligonucleotides can activate various immune subsets, including natural killer cells (which produce interferon  $\gamma$  and have  
15 cytolytic activity) and macrophages (Wooldrige et al Vol 89 (no. 8), 1977). Although other unmethylated CpG containing sequences not having this consensus sequence have now been shown to be immunomodulatory.

The present invention provides an improved vaccine formulation comprising a CpG  
20 oligonucleotide and an HIV antigen selected from the group HIV gp160, HIV gp120; an HIV Nef protein or derivative thereof linked to either (i) a fusion partner or (ii) an HIV Tat protein or derivative thereof; or an HIV Tat protein or derivative thereof linked to either (i) a fusion protein or (ii) an HIV Nef protein or derivative thereof; or an HIV Nef protein or derivative thereof linked to an HIV Tat protein or  
25 derivative thereof and a fusion partner.

By 'fusion partner' is meant any protein sequence that is not Tat or Nef. Preferably the fusion partner is protein D or its lipidated derivative Lipoprotein D, from *Haemophilus influenzae* B. In particular, it is preferred that the N-terminal third,  
30 i.e. approximately the first 100-130 amino acids are utilised. This is represented herein as Lipo D 1/3. In a preferred embodiment of the invention the Nef protein



or derivative thereof may be linked to the Tat protein or derivative thereof. Such Nef-Tat fusions may optionally also be linked to a fusion partner, such as protein D.

If present, the fusion partner is normally linked to the N-terminus of the Nef or Tat  
5 protein.

Derivatives encompassed within the present invention include molecules with a C terminal Histidine tail, which preferably comprises between 5-10 Histidine residues. Generally, a histidine tail containing n residues is represented herein as His (n).

10 The presence of a histidine (or 'His') tail aids purification. More specifically, the invention provides vaccines comprising proteins with the following structure:

	Lipo D 1/3	-	Nef	-	His (6)
15	Lipo D 1/3	-	Nef-Tat	-	His (6)
	Prot D 1/3	-	Nef	-	His (6)
	Prot D 1/3	-	Nef-Tat	-	His (6)
20			Nef-Tat		His (6)

In a preferred embodiment the proteins are expressed with a Histidine tail comprising between 5 to 10 and preferably six Histidine residues. These are  
25 advantageous in aiding purification. Separate expression, in yeast (*Saccharomyces cerevisiae*), of Nef (Macreadie IG et al 1993, *Yeast* 9 (6) 565-573) and Tat (Braddock M et al, 1989, *Cell* 58 (2) 269-79) has already been reported. Nef protein only is myristilated. The present invention also provides vaccines comprising Nef and/or Tat separately. The DNA and amino acid sequences of  
30 representative Nef-His, Tat-His and of Nef-Tat-His fusion proteins as set forth in WO99/16884.

Particularly preferred vaccines according to the invention comprise an HIV fusion protein such as Nef-Tat and optionally an additional HIV protein. Particularly preferred in a vaccine formulation according to the invention is a combination of an HIV fusion protein plus gp120, more particularly Nef-Tat with gp120.

5

Derivatives encompassed within the present invention also include mutated proteins. The term 'mutated' is used herein to mean a molecule which has undergone deletion, addition or substitution of one or more amino acids using well known techniques for site directed mutagenesis or any other conventional method.

10

Vaccine preparation is generally described in Vaccine Design - The subunit and adjuvant approach (Ed. Powell and Newman) Pharmaceutical Biotechnology Vol. 6 Plenum Press 1995. Encapsulation within liposomes is described by Fullerton, US Patent 4,235,877.

15

The preferred oligonucleotides preferably contain two or more CpG motifs separated by six or more nucleotides. The oligonucleotides of the present invention are typically deoxynucleotides. In a preferred embodiment the internucleotide in the oligonucleotide is phosphorodithioate, or more preferably a phosphorodithioate bond, although phosphodiester and other internucleotide bonds are within the scope of the invention including oligonucleotides with mixed internucleotide linkages.

20

Preferred oligonucleotides have the following sequences:

Oligo (internal designation*)	5'-SEQUENCE-3'	CpG	Thio
WD1001	TCC ATG ACG TTC CTG ACG TT	+	+
WD1002	TCT CCC AGC GTG CGC CAT	+	+
WD1003	ACC GAT AAC GTT GCC GGT GAC G	+	-
WD1004	G*G*G GTC AAC GTT GAG* G*G*G* G*G	+	Mix
WD1005	TCC ATG AGC TTC CTG AGC TT	-	+

WD1006	TCC ATG ACG TTC CTG ACG TT	+	-
WD1007	ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG	+	+
	TCG TCG TTT TGT CGT TTT GTC GTT	+	+

\* alternatively referred to as WD001-WD007

In the above table a + in the Thio column indicates the presence of a thioate modification. 'Mix' indicates a mixture of thioate modification and sequence without thioate modification (the asterisks indicate the linkages with a thioate modification). A - in the Thio column indicates absence of a thioate modification. A + in the CpG column indicates a the presence of a CpG motif and a - in the CpG column indicates absence of a CpG motif. For example WD1005 contains a GpC rather than a CpG motif, thus it is marked with a - in the CpG column of the table. WD1007 contains a palindromic motif (GACGTC) as well as other non-palindromic CpG sequences. This is also within the scope of a CpG oligonucleotide as the term is used in the present application.

The CpG oligonucleotides utilised in the present invention may be synthesized by any method known in the art (eg EP 0 468 520). Conveniently, such oligonucleotides may be synthesized utilising an automated synthesizer. Methods for producing phosphorothioate oligonucleotides or phosphorodithioate are described in US 5,666,153, US 5,278,302 and WO 95/26204.

The amount of protein in each vaccine dose is selected as an amount which induces an immunoprotective response without significant, adverse side effects in typical vaccinees. Such amount will vary depending upon which specific immunogen is employed and how it is presented. Generally, it is expected that each dose will comprise 1-1000 µg of protein, preferably 2-500 µg, most preferably 5-250 µg. An optimal amount for a particular vaccine can be ascertained by standard studies involving observation of appropriate immune responses in subjects. Following an initial vaccination, subjects may receive one or several booster immunisations adequately spaced.

It is also possible to pre-administer the CpG oligonucleotide shortly prior to vaccination with the HIV antigen, for example 1 day before.

Accordingly, according to another aspect of the invention there is provided a method of treating a human in need thereof either with a vaccine composition comprising an HIV antigen and a CpG oligonucleotide (as hereinabove defined) or with the CpG oligonucleotide followed after a suitable time by the HIV antigen.

There is also provided a kit comprising effective amounts of a CpG oligonucleotide-containing formulation for use as a priming formulation for pre-administration to human patients and an HIV antigen for injection at some suitable time later, as described hereinabove.

Preferred CpG oligonucleotides are those indicated in the table hereinabove.

Suitably the CpG will be present in the range 10 µg per dose to 2000 µg, preferably 50-1000 µg, such as about 50 or about 500 or about 1000 µg per dose.

Suitably the vaccine used in the present invention may comprise a carrier such as an aluminium salt, eg aluminium hydroxide (Al(OH)<sub>3</sub>), aluminium phosphate or aluminium phosphate sulfate (alum), or a non-toxic oil in water emulsion or a mixture thereof.

If an aluminium salt (preferably aluminium hydroxide) is used as a carrier it is generally present in the range of 50 to 100 µg, preferably 100 to 500 µg per dose.

Non-toxic oil in water emulsions preferably contain a non-toxic oil, eg squalene and an emulsifier such as (polysorbitan monoleate) Tween 80, in an aqueous carrier such as phosphate buffered saline.

If desired the vaccine used in the present invention may comprise an additional adjuvant, preferably a saponin adjuvant such as QS21 as described for example in

WO 9517210, optionally in the presence of a sterol, such as cholesterol as described for example in PCT/EP96/01464. The vaccine of the invention may also comprise monophosphoryl lipid A and derivatives thereof known in the art. A preferred derivative is 3 de-O-acylated monophosphoryl lipid A, described in British Patent

5 No. GB 2 220 211.

Accordingly vaccine formulations of the present invention may additionally comprise other pharmaceutical excipients or immunostimulants. In a preferred embodiment the vaccine formulation additionally comprises an aluminium salt,

10 preferably aluminium hydroxide. In a further embodiment a saponin adjuvant may also be included such as QS21 (Aquila).

The present invention will now be described with reference to the following examples:

## EXAMPLES

### Example 1 - IMMUNOGENICITY STUDIES USING HIV-1 GP120

#### 5 FORMULATED WITH CPG OR CPG/ALUM

#### Evaluation of CpG and CpG/alum in rhesus monkeys

##### *Experiment outline*

An immunogenicity study was conducted to evaluate the adjuvant effect of CpG in non-human primates. Groups of five monkeys were immunized twice with HIV-  
 10 1<sub>W6.1D</sub> gp120 in combination with CpG or CpG/alum. After the second immunization the immune response of the animals was assessed. Antibodies to gp120 and lymphoproliferative as well as cytokine responses were evaluated.

#### Groups of monkeys

15

Group	antigen	Adjuvant
1	HIV-1 <sub>W6.1</sub> gp120	CpG/alum
2	HIV-1 <sub>W6.1</sub> gp120	CpG

#### Formulation

##### *Component batches used:*

20

COMPONENT	BRAND	BATCH NUMBER	CONCENTRATION (MG/ML)	BUFFER
Gp120*		47/025	0.456	PBS
Al(OH) <sub>3</sub>	Superfos	96A0089	10.380	H <sub>2</sub> O
CpG		WD001	5	H <sub>2</sub> O

\* gp120 of HIV-1 clone W6.1D (Groenink et al., 1991)

**Formulation process:**

Formulations were prepared one day before each injection. All incubations were carried out at room temperature with agitation.

5

**CpG/alum group 1 (500µl/dose)**

Gp120 (50µg) was adsorbed on 500µg of Al(OH)<sub>3</sub> for 1 hour. The formulation was buffered with a 10-fold concentrated PO<sub>4</sub>/NaCl pH 6.8 solution before addition of  
10 500µg of WD001. After 15 min, 50 µg/ml of thiomersal was added as preservative.

**CpG group 2 (500µl/dose)**

Gp120 (50µg) was diluted in PO<sub>4</sub>/NaCl buffer pH 6.8 before addition of 500µg of  
15 CpG WD001. After 15 min, 50 µg/ml of thiomersal was added as preservative.

**Immunological methods**

Five rhesus monkeys (*Macaca mulatta*). per group were immunized twice  
20 intramuscularly with 500 µl of vaccine at a four-week-interval. Sera and peripheral blood mononuclear cells (PBMC) were taken at several occasions.

Gp120-specific antibodies in sera of immunized monkeys were determined in an ELISA as described in Mooij et al., 1998. Briefly, ELISA plates were coated with 1  
25 µg/ml gp120, saturated with BSA and subjected to incubation with rhesus monkey sera followed by α-human biotinylated antibody. Finally, HRP-streptavidin was added and quantified in a color reaction using OPD as substrate.

Lymphoproliferation was assessed by using density gradient-purified PBMC from  
30 immunized rhesus monkeys. Cells were seeded in quadruplicates at 1x10<sup>5</sup> in 100 µl RPMI/5 % FCS per well in round bottom 96 well plates. Then another 100 µl of

medium alone or containing soluble gp120 (10  $\mu$ g/ml) were added and parallel cultures were incubated for 48 hrs. Thereafter, 100  $\mu$ l culture supernatant were replaced by fresh medium containing 1  $\mu$ Ci [ $^3$ H]-thymidine. After 16 hrs cells were harvested onto filter plates and incorporated radioactivity was determined in a  $\beta$ -counter. Results are expressed in cpm and in stimulation indices (SI, = cpm antigen-containing cultures/cpm medium alone cultures), SI greater than 3 are considered as a positive response.

Flat bottom 96 well plates were prepared by coating an IFN- $\gamma$ -specific capture antibody in 50  $\mu$ l PBS for 4 hrs at 37  $^{\circ}$ C. The plates were washed three times and PBMC were seeded similar to lymphoproliferation assays. After 48 hrs of culture the plates were washed thrice with PBS/0.05 % Tween 20 and 50  $\mu$ l of biotinylated secondary IFN- $\gamma$ -specific antibody diluted in PBS/Tween/1 % FCS were added for 2 hrs. The plates were washed again and a gold-conjugated  $\alpha$ -biotin antibody was incubated for 1 hr. After additional washings the ELIspots were visualized by using a silver enhancing kit (50  $\mu$ l per well). The reaction was stopped after approx. 30 min by adding deionized water. Cytokine-secreting cells were enumerated by microscopic examination.

## 20 Results

Analysis of gp120-specific antibodies in sera of the monkeys revealed that all animals in the two groups had acquired specific immune responses (Table 1 and Figure 1). Some low responses were detectable in two animals already after one immunization. After the second immunization all animals exhibited high titers of gp120-specific antibodies.



Table 1

		d - 12	average	14 d post I	average	7 d post II	average	28 d post II	average
Group 1	N5V	<100		188		6445		2014	
	1055	<100		50		4699		1484	
	R441	<100	<100	50	<50	2942	5530	827	2504
	BXJ	<100		50		5808		4013	
	CPJ	<100		50		7758		4184	
Group 2	NJ5	<100		50		734		729	
	X006	<100		50		4182		1171	
	1041	<100	<100	50	<50	2834	3358	400	963
	R361	<100		258		5227		1821	
	BBE	<100		50		3811		693	

- 5 Induction of specific lymphoproliferation by immunization with gp120 in combination with CpG or CpG/alum was evaluated before immunization and 6 days post secondary immunization. All 10 animals did not exhibit any specific lymphoproliferation (SI > 3) at the study start (data not shown). In contrast, all animals in group 1 possessed strong lymphoproliferative responses 6 days post boost immunization (Figure 2). Similarly, all except one animal in group 2 showed specific lymphoproliferation.

The presence of gp120-specific IFN- $\gamma$ -secreting cells was investigated in all monkeys before immunization and 6 days after the second dose. IFN- $\gamma$ -secreting cells could not be evaluated from pre-immunization samples due to technical difficulties. However, such cells were detectable after secondary immunization (Figure 3). All animals in both groups exhibited an antigen-specific response.

### Conclusions


- 20 Immunization with HIV-1 gp120 in combination with CpG or CpG/alum induces immune responses in non-human primates. These responses include gp120-specific antibodies, lymphoproliferation and IFN- $\gamma$ -secreting cells.

**Example 2 - NEF-TAT AND MUTATED NEF-TAT FORMULATION**

The antigens prepared according to WO99/16884 were dialysed against NaCl 150mM to eliminate the  $\text{PO}_4$  ions that inhibit the adsorption of gp120 on  $\text{Al}(\text{OH})_3$ .

5,

The formulation sequence is as described below:



- 20 The antigens were incubated with the CpG before adsorption on  $\text{Al}(\text{OH})_3$  to favour a potential interaction between the His tail of the Nef-Tat and Nef antigens and the oligonucleotide.

**Formulation Characterisation**

- 25 The formulation was centrifuged and the supernatant was analysed by SDS-PAGE (Novex 4-20%, reducing conditions) to check the adsorption of HIV Nef-Tat. The gel was stained with silver. Gel analysis of the formulation revealed that most of the Nef or Nef-Tat protein was adsorbed onto alum.

30

**Example 3 - IMMUNIZATION AND SHIV CHALLENGE EXPERIMENT IN RHESUS MONKEYS.*****Vaccine preparation***

- 5 A vaccine formulation was prepared in accordance with the invention comprising a Nef-Tat fusion protein and gp120 together with an oligonucleotide containing unmethylated CpG dinucleotide motifs and aluminium hydroxide as carrier.

Preparation of CpG oligonucleotide solution: CpG dry powder is dissolved in H<sub>2</sub>O to  
10 give a solution of 5 mg/ml CpG.

**Preparation of CpG formulation.**

The 3 antigens were dialyzed against NaCl 150 mM to eliminate the phosphate ions that inhibit the adsorption of gp120 on aluminium hydroxide.

15

- The antigens diluted in H<sub>2</sub>O (100 µg gp120, 20 µg NefTat and 20 µg SIV Nef) were incubated with the CpG solution (500 µg CpG) for 30 min before adsorption on Al(OH)<sub>3</sub> to favor a potential interaction between the His tail of NefTat and Nef antigens and the oligonucleotide (stronger immunostimulatory effect of CpG  
20 described when bound to the antigen compared to free CpG). Then Al(OH)<sub>3</sub> (500 µg), 10 fold concentrated NaCl and 1 µg/ml thiomersal as preservative were consecutively added at 5 min intervals.

All incubations were carried out at room temperature with agitation.

25

***Study***

Four rhesus monkeys were immunized intramuscularly at 0, 1 and 3 months with the vaccine composition.

One month after the last immunization all animals were challenged with a pathogenic SHIV (strain 89.6p). From the week of challenge (wk16) blood samples were taken periodically at the indicated time points to determine the % of CD4-positive cells among peripheral blood mononuclear cells by FACS analysis (Figure 4) and the  
5 concentration of RNA viral genomes in the plasma by bDNA assay (Figure 5).

### ***Results***

All animals became infected after challenge with SHIV<sub>89.6p</sub>.

- 10 CD4-positive cells declined after challenge. All four animals exhibited a slight decrease in CD4-positive cells and recovered to baseline levels over time (Figure 4). In contrast, three out of four animals in an adjuvant alone control group exhibited a drop of CD4-positive cells below 5 % and never recovered.
- 15 Virus load data are almost the inverse of CD4 data (Figure 5). Virus load decreased in the animals that had received NefTat + gp120 formulated with CpG containing oligonucleotide, but remained high in three out of four control animals which received only an adjuvant.
- 20 At week 44 2/4 animals in a control group treated with only an adjuvant had to be euthanized due to AIDS-like symptoms.

### ***Conclusions***

- The combination of gp120 and NefTat (in the presence of SIV Nef) with CpG  
25 adjuvant prevents the loss of CD4-positive cells and reduces the virus load in animals infected with pathogenic SHIV<sub>89.6p</sub>.

**Figure legends**

Figure 1: HIV-1 gp120-specific antibody responses in immunized rhesus monkeys. Specific antibodies were evaluated in an ELISA. Individual values from multiple  
5 time points for each animal are shown in Table 1, and group averages are shown in Table 1 and as a graphic in figure 1.

Figure 2: Gp120-specific lymphoproliferation in immunized rhesus monkeys 6 days post second immunization. PBMC were stimulated with gp120 antigen and  
10 lymphoproliferative responses were measured by <sup>3</sup>H-thymidine incorporation. Results are expressed in cpm and as SI.

Figure 3: Gp120-specific IFN- $\gamma$ -secreting cells from immunized rhesus monkeys. IFN- $\gamma$ -secreting cells were visualized by the ELISpot method. Cytokine-secreting  
15 cells resulting in a colored spot were enumerated by microscopic examination and results are expressed semi-quantitatively (- = 0-5, + = 5-15, ++ = 15-35, +++ = 35-50, ++++ = >50).

Figures 4 and 5: CD4 cell count and viral load data from Example 3 in which  
20 rhesus monkeys were immunised with Nef-Tat + gp120 formulated with CpG and challenged with pathogenic SHIV.

**References**

- Groenink, M., Fouchier, R.A., De Goede, R.E., et al. (1991). Phenotypic  
25 heterogeneity in a panel of infectious molecular human immunodeficiency virus type 1 clones derived from a single individual. *J. Virol.* 65:1968-1975.
- Mooij, P., Van der Kolk, M., Bogers, M.J.M., et al. (1998). A clinically  
relevant HIV-1 subunit vaccine protects rhesus macaques from in vivo passaged  
30 simian-human immunodeficiency virus infection. *AIDS* 12:F15-F22.

### Claims

1. A vaccine formulation comprising a HIV antigen and an immunostimulatory CpG oligonucleotide.
- 5 2. A vaccine as claimed in claim 1 wherein the antigen is selected from the group, gp120, gp160, Nef, Tat, Nef-Tat fusions or immunologically equivalent derivatives.
3. A vaccine as claimed in claim 1 or 2 wherein the antigen is fused to Protein D,  
10 or lipoprotein D or a fragment thereof, from Haemophilus influenzae.
4. A vaccine as claimed in any one of claims 1 to 3 wherein the antigen comprises a Nef-Tat fusion protein
- 15 5. A vaccine as claimed in claim 4 further comprising HIV gp120.
6. A vaccine formulation as claimed herein additionally comprising an aluminium salt or a saponin adjuvant.
- 20 7. A vaccine as claimed herein wherein the oligonucleotide comprises two CpG dinucleotides.
8. A vaccine as claimed herein wherein the CpG oligonucleotide is between 15-45 nucleotides in length.
- 25 9. A vaccine as claimed herein wherein the CpG oligonucleotide comprises at least one phosphorothioate internucleotide bond.
10. A vaccine as claimed herein wherein the oligonucleotide is selected from the  
30 group:

Oligo (internal designation)	5'-SEQUENCE-3'	CpG	Thio
WD1001	TCC ATG ACG TTC CTG ACG TT	+	+
WD1002	TCT CCC AGC GTG CGC CAT	+	+
WD1003	ACC GAT AAC GTT GCC GGT GAC G	+	-
WD1004	G*G*G GTC AAC GTT GAG* G*G*G* G*G	+	Mix
WD1005	TCC ATG AGC TTC CTG AGC TT	-	+
WD1006	TCC ATG ACG TTC CTG ACG TT	+	-
WD1007	ACC GAT GAC GTC GCC GGT GAC GGC ACC ACG	+	+
	TCG TCG TTT TGT CGT TTT GTC GTT	+	+

11. A method for the prevention or amelioration of HIV infection a patient, comprising administering an effective amount of a vaccine of claims 1 to 10 to patient.

5


12. A vaccine as claimed herein for use as a medicament.

13. A method of producing a vaccine as claimed in any of claim 1 to 8 comprising admixing an HIV antigen and a CpG immunostimulatory oligonucleotide.

10

14. A method for the prevention or amelioration of HIV infection in a patient, comprising administering an effective amount of a CpG oligonucleotide followed after a suitable time by an effective amount of an HIV antigen.

1/5

**Fig. 1**



2/5




Fig. 2b




Fig. 2d




Fig. 2a




Fig. 2c

Fig. 2

3/5

Group 1					
	N5V	1055	R441	BXJ	CPJ
gp120	+	++	++	+	+++
Unstimulated cells	-	-	-	-	-
					5/5

Group 2					
	NJ5	X006	1041	R361	BBE
gp120	+++	+++	++	+++	+
Unstimulated cells	-	+	+	+	-
					5/5

Fig. 3

4/5




Fig. 4

5/5




Fig. 5